

get to know **KRAZATI**[®]

(adagrasib) | 200 mg TABLETS

GET TO KNOW KRAZATI TO HELP MAKE THE NEXT MOVE AGAINST KRAS G12C-POSITIVE ADVANCED NSCLC

How KRAZATI is designed for my specific type of cancer

Q: What is KRAZATI?

A: Designed to target your specific type of cancer, KRAZATI helps block the abnormal KRAS G12C protein, targeting it in a way chemotherapy and immunotherapy do not.

KRAZATI is for adults with NSCLC that has spread to other parts of the body or cannot be removed by surgery and have received prior systemic (or patient level language equivalent) treatment.

Before prescribing KRAZATI, your doctor will do a test, called a biomarker test, to see if you have KRAS G12C-positive NSCLC.

Q: What is a mutation?

A: A mutation is a change to one or more of a cell's genes. Usually, our immune system finds and destroys cells with mutated genes. Sometimes, though, a mutation like KRAS G12C can cause cancer cells to grow and spread quickly and avoid the immune system.

Q: What is a biomarker?

A: A biomarker is a biological feature, such as the KRAS G12C mutation, that can be used to identify different cancer types. Samples taken from the cancer are used to find biomarkers. Once you have your biomarker test results, you and your doctor can plan the appropriate treatment for your type of NSCLC.



Important Facts About **KRAZATI**[®] (adagrasib)

This is a summary of important information that you need to know about KRAZATI. Your healthcare team can work with you to help answer any questions you may have. **Keep this document in a safe place so you can refer to it before you start taking it and during your treatment.**

Look out for the following icons as you read:



Talk to your healthcare team



Call a healthcare provider right away



Helpful information to remember

Please see additional Important Facts about KRAZATI throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.

Q: How does KRAZATI® work?

A: KRAZATI helps fight your cancer in a way chemotherapy and immunotherapy do not. It targets and helps block the KRAS G12C mutation that can cause tumors to grow and spread throughout the body.



Fitting KRAZATI into my daily routine

Q: How often do I need to take KRAZATI?

A: Take KRAZATI exactly as your doctor tells you to take it. The recommended dosage is 600 mg twice daily. Do not change your dose or stop taking KRAZATI unless your healthcare provider tells you to. Your healthcare provider may change your dose, or temporarily or permanently stop treatment with KRAZATI if you develop certain side effects.

Q: Do I need to take KRAZATI at the same time every day?

A: Take your prescribed dose of KRAZATI at the same time every day.

Q: How do I take KRAZATI?

A: You can take KRAZATI with or without food. Swallow each tablet whole, and don't chew, crush, or split the tablets.



Important Facts About KRAZATI (cont'd)

What is KRAZATI?

KRAZATI is an oral prescription medicine used to treat:

Your healthcare team will perform a test to make sure that KRAZATI is right for you.

It is not known if KRAZATI is safe and effective in children.

Talk to your healthcare team for more information about what it means to have an abnormal KRAS G12C gene.

Non-small cell lung cancer (NSCLC) in adults who have:

- ✓ cancer that has spread to other parts of the body or cannot be removed by surgery, **AND**
- ✓ a tumor that has an abnormal KRAS G12C gene, **AND**
- ✓ received at least one prior medicine that works throughout your whole body (a systemic treatment)

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.

Q: What if I miss a dose of KRAZATI®?

A: If you miss a dose of KRAZATI, take it as soon as you remember. However, if it has been more than 4 hours since the time of the missed dose, skip that dose, and take the next dose of KRAZATI at the usual time. Don't take 2 doses at the same time to make up for a missed dose.

Q: What if I vomit right after taking KRAZATI?

A: If you vomit after taking KRAZATI, do not take an extra dose. Take the next dose at your next scheduled time.

Q: What if I take other medications?

A: Talk to your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. KRAZATI can affect the way other medicines work, and other medicines can affect how KRAZATI works.

Q: What if I can't swallow KRAZATI whole?

A: Let your doctor know if you have any trouble swallowing KRAZATI whole.



The side effects I may see with KRAZATI

Q: What are the possible side effects I may experience on KRAZATI?

A: There is a possibility you will experience side effects while on KRAZATI. KRAZATI may cause serious side effects, including stomach or intestinal (gastrointestinal). Changes in the electrical activity of your heart called QTc prolongation with your liver, and with your lungs or breathing. The most common side effects of KRAZATI include: nausea, diarrhea, vomiting, tiredness, muscle and bone pain, kidney problems, swelling, decreased appetite, and trouble breathing.

Certain abnormal blood test results are common during treatment with KRAZATI. Your healthcare provider will monitor you for abnormal blood tests and treat you if needed.

KRAZATI may cause fertility problems in males and females and may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.

Tips for managing the more common side effects



Q: What do I do if I experience side effects while taking KRAZATI®?

A: It's important to talk to your doctor right away if you experience any side effects.

Q: What are some tips to manage side effects?

A: KRAZATI can cause serious side effects. It's important to talk to your doctor right away if you experience any side effects. Side effects can vary from person to person. Here are some ways that may help you manage them:

Diarrhea

- Drinking fluids may help you stay hydrated
- Eat smaller meals with foods that might be easy on your stomach. Foods that are high in potassium and sodium may help replace minerals that your body has lost
- Discuss other possible treatments with your doctor

Nausea and vomiting

- Drink plenty of fluids—clear liquids and ice cold beverages
- Eat smaller, more frequent meals. Nausea may be worse on an empty stomach
- Pay attention to which foods bother your stomach and try to avoid them (ie, greasy, fried, sweet, or spicy foods)
- Talk with your doctor about possibly taking anti-nausea medicine

Tiredness/fatigue

- Track your energy level throughout the day, and have it on hand to discuss with your doctor
- Balance rest and activity, prioritizing your most necessary activities for when you may have the most energy
- Staying hydrated and eating foods high in protein may increase your energy levels
- If feelings/emotions are disturbing your sleep, talk to your doctor about a referral to a specialist

Important Facts About KRAZATI (cont'd)

Before taking KRAZATI, what should I discuss with my healthcare team?

Talk to your healthcare team about all of your medical conditions, including if you have:

- any heart problems, including heart failure or if you were born with long QT syndrome
- liver problems

You should not take KRAZATI if you were born with long QT syndrome or if you currently have QTc prolongation (which means your heart takes longer than usual to recharge between beats).

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.

Where I can go for support and information



Q: Who can help me with insurance coverage and questions about paying for KRAZATI®?

A: BMS Access Support provides resources and assistance to:

- Help you understand your insurance coverage
- Support and educate you throughout each stage of your treatment
- Inform you about financial support options, including co-pay assistance for eligible commercially-insured patients

For more information, ask your doctor, visit www.BMSAccessSupport.com, or call BMS Access Support at **1-800-861-0048**, 8 AM to 8 PM ET, Monday-Friday.

A: KRAZATI is only available with a prescription from your doctor. Once prescribed, KRAZATI is available through the pharmacy at your doctor's office or through a specialty pharmacy that will fill the prescription by mail. KRAZATI is not available at your local retail pharmacy.

DISCOVER MORE
SUPPORT AND
RESOURCES



Important Facts About KRAZATI (cont'd)

Talk to your healthcare team about all the medicines you take, including:

- prescription medicines
- over-the-counter medicines
- vitamins
- herbal supplements

Taking KRAZATI with certain other medicines may affect how KRAZATI or your other medicines work. It may also increase your chance of getting side effects from the medicines you take.

Keep a list of the medicines you take and show it to your healthcare team and pharmacist before you start taking KRAZATI or any other new medicine.

Talk to your healthcare team about anything you might be unsure of before starting treatment. There may be other topics that are important to discuss with your healthcare team that aren't in this document.

Before taking KRAZATI, what should I discuss with my healthcare team about pregnancy, breastfeeding, and fertility?

Talk to your healthcare team if:

You are pregnant or plan to become pregnant: It is not known if KRAZATI can harm your unborn baby.

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.

Important Facts About KRAZATI® (cont'd)

Before taking KRAZATI, what should I discuss with my healthcare team about pregnancy, breastfeeding, and fertility? (cont'd)

 **Talk to your healthcare team if:**

-  **You are breastfeeding' or plan to breastfeed:** It is not known if KRAZATI passes into your breast milk. Do not breastfeed during your treatment and for 1 week after your last dose of KRAZATI.

KRAZATI may cause fertility problems in all genders and may affect your ability to have children.

 **Talk to your healthcare team if having fertility problems is a concern for you.**

What are the serious side effects of KRAZATI?

KRAZATI can cause serious side effects. A serious side effect is a side effect that may require medical treatment or hospitalization, cause permanent damage, or be life-threatening or sometimes even fatal. Your healthcare team may lower your dose, pause, or stop treatment with KRAZATI if you get certain side effects. The serious side effects of KRAZATI may include:

Problems in your stomach and intestines (gastrointestinal problems). Problems in your stomach and intestines can be severe when taking KRAZATI. These problems may include:

- severe nausea, vomiting, or diarrhea
- inflammation in your colon (colitis)
- bleeding in your stomach or intestines (gastrointestinal bleeding)
- certain parts of your stomach or intestines becoming narrower than normal (gastrointestinal stenosis)
- a blockage in your stomach or intestines (bowel obstruction)

 **Call a healthcare provider right away** if you have any of the following signs or symptoms of problems in your stomach and intestines:

- vomiting six or more times (separated by 5 minutes) in 24 hours
- dark-colored stool or stool with blood in it
- vomit with blood in it or vomit that looks like coffee grounds
- diarrhea where you have at least 7 more stools per day than usual and have trouble holding in your bowel movements (fecal incontinence)
- nausea that prevents you from eating and drinking enough
- feeling dizzy, lightheaded, or weak
- pain in your stomach (abdominal) area
- shortness of breath
- trouble urinating or dark urine
- feeling like your heart is beating faster than normal or fluttering (rapid pulse)

Your healthcare team may prescribe you a medicine (such as an anti-diarrheal or anti-nausea medicine) to help treat possible side effects related to your stomach and intestines.

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.



Important Facts About KRAZATI® (cont'd)

What are the serious side effects of KRAZATI? (cont'd)

Changes to the way your heart beats (called QTc prolongation). Taking KRAZATI may cause changes to the way your heart beats, including causing your heart to take longer to recharge between beats (this is called QTc prolongation). QTc prolongation may increase your chance of developing an irregular heartbeat that may be life-threatening and lead to sudden death. For example, it may cause a particular kind of very fast heartbeat called Torsades de Pointes.

Before and during your treatment with KRAZATI, your healthcare team may:

- give you a test called an electrocardiogram (ECG) to measure the electrical activity of your heart
- give you blood tests to measure how much of certain kinds of minerals (electrolytes) like potassium and magnesium are in your blood

Your healthcare team is more likely to give you the above tests if you have heart failure, a slow heart rate, abnormal levels of electrolytes in your blood, or are taking another medicine that may lead to QTc prolongation.

 **Call a healthcare provider** if you have any of the following signs or symptoms:

- feeling dizzy, lightheaded, or weak
- an abnormal heartbeat (arrhythmia) such as your heart beating faster than usual, slower than usual, or beating in an irregular pattern

Problems with your liver. It is common to get abnormal results on liver blood tests when you take KRAZATI. These abnormal results may sometimes indicate severe problems with your liver. Your healthcare team will do blood tests before and during your treatment with KRAZATI to check how your liver is working.

 **Call a healthcare provider** if you have any of these signs or symptoms:

- yellowing of your skin or the white parts of your eyes (jaundice)
- unexplained nausea or vomiting
- dark-colored urine
- bleeding or bruising more easily than normal
- light-colored stool
- losing your appetite
- feeling tired or weak
- pain, aching, or tenderness on the right side of your stomach (abdominal) area

Problems with your lungs or breathing (lung inflammation or pneumonitis): KRAZATI may cause inflammation in your lungs, which may be severe or serious and can lead to death.

 **Call a healthcare provider or get emergency medical help right away** if you have any of these signs or symptoms:

- new or worsening shortness of breath
- fever
- new or worsening cough

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.



Important Facts About KRAZATI® (cont'd)



What are the most common side effects of KRAZATI?

The most common side effects of KRAZATI when it is taken on its own for **non-small cell lung cancer (NSCLC)** include:

- nausea
- diarrhea
- vomiting
- feeling tired
- pain in your muscles and bones
- kidney problems
- swelling in any part of your body
- feeling less hungry than usual (decreased appetite)
- trouble breathing

It is also common to get abnormal results on certain blood tests while you are taking KRAZATI. While you are taking KRAZATI, your healthcare team will give you blood tests to monitor you for abnormal blood test results and will treat you if needed.

The common and serious side effects mentioned in this document are not all the possible side effects of KRAZATI.

 **Talk to your healthcare team for more information or advice about side effects.** You are encouraged to report side effects to the FDA at www.fda.gov/medwatch or calling 1-800-FDA-1088.

What does KRAZATI look like and what are its ingredients?

KRAZATI comes as a white or off-white oval-shaped tablet, each containing a 200mg dose of its active ingredient. Each tablet has the number “200” stamped into the surface on one side, and a stylized “M” stamped into the other side.



Note: Image does not represent actual tablet sizes

Ingredients:

- **Active:** adagrasib
- **Inactive:** colloidal silicon dioxide, crospovidone, magnesium stearate (vegetable sourced), mannitol, microcrystalline cellulose, hypromellose, maltodextrin, medium chain triglycerides (vegetable sourced), polydextrose, talc, and titanium dioxide

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.

Important Facts About KRAZATI® (cont'd)



How should I take KRAZATI?

Take KRAZATI exactly as your healthcare team tells you. Do not change how much you take or stop taking KRAZATI unless a healthcare provider tells you to.

The recommended dose of KRAZATI is three tablets (a 600mg dose) by mouth twice a day, at about the same time each day. Your healthcare team may need to change how much KRAZATI you take, or pause or stop your treatment with KRAZATI, if you have certain side effects.

- ✓ **Do** take KRAZATI by mouth (orally) twice a day, around the same time each day
- ✓ **Do** take KRAZATI with or without food
- ✓ **Do** swallow KRAZATI tablets **whole**
- ✗ **Do not** chew, crush or split tablets
- ✗ **Do not** take broken or damaged tablets

What if I vomit or miss a dose of KRAZATI?

Do not take an extra dose.

- **If you vomit after taking a dose** of KRAZATI, do not take an extra dose. Take your next dose at your next scheduled time.
- **If you miss a dose** of KRAZATI and it has been:
 - **Less than four hours** since you were supposed to take it, take the dose.
 - **More than four hours** since you were supposed to take your dose, skip that dose. Just take your next dose of KRAZATI at your next scheduled time. Do not take two doses at the same time to make up for a missed dose.



Setting an alarm clock or a calendar notification is a great way to help you remember to take your medication.

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.

Important Facts About KRAZATI® (cont'd)

KRAZATI®
(adagrasib) | 200 mg
TABLETS

What if I take too much KRAZATI?

 **Call a healthcare provider or Poison Health Line at (1-800-222-1222), or go to the nearest hospital emergency room right away.**

How should I store KRAZATI?

Keep your KRAZATI tablets in their original containers. These containers are child-resistant and include a packet of desiccant (drying agent) that keeps tablets dry.

- **Do not** remove the packet of desiccant (drying agent) from the container.
- **Do not** eat or swallow the desiccant.

Store tablets at **room temperature** between 68°F to 77°F (20°C to 25°C).

Keep KRAZATI and all medicines out of reach of children.

What else should I know about KRAZATI?

Do not use KRAZATI for any condition other than what your healthcare team has prescribed it for. Healthcare teams may sometimes prescribe KRAZATI for purposes other than those mentioned in this document.

Do not give KRAZATI to other people, even if they have the same symptoms as you. It may harm them.

Ask your healthcare team for the information about KRAZATI that is written for healthcare providers if you would like.

 For more information, please see the accompanying US Full Prescribing Information and Patient Information for KRAZATI. Talk to your healthcare team for more information about this medication.

Please see additional Important Safety Information throughout and the Patient Information in the accompanying Full Prescribing Information in pocket.

References:

1. KRAZATI [package insert]. Princeton, NJ: Bristol-Myers Squibb Company; 2024. 2. Docetaxel [package insert]. Pfizer Inc. 2022. 3. Pakkala S, Ramalingam SS. Personalized therapy for lung cancer: striking a moving target. *JCI Insight*. 2018. doi:10.1172/jci.insight.120858 4. National Cancer Institute. Dictionary of cancer terms. National Cancer Institute website. Accessed October 29, 2025. <https://www.cancer.gov/publications/dictionaries/cancer-terms/> 5. Janne PA, Riely GJ, Gadgeel SM, et al. Adagrasib in non-small-cell lung cancer harboring a KRAS^{G12C} mutation. *N Engl J Med*. 2022;387(2):120-131. 6. National Cancer Institute. Diarrhea and cancer treatment. National Cancer Institute website. Updated May 16, 2025. Accessed August 27, 2025. <https://www.cancer.gov/about-cancer/treatment/side-effects/diarrhea>. 7. National Cancer Institute. Nausea and vomiting and cancer treatment. National Cancer Institute website. Updated May 9, 2025. Accessed August 27, 2025. <https://www.cancer.gov/about-cancer/treatment/side-effects/nausea-vomiting>. 8. American Cancer Society. Managing nausea and vomiting at home. American Cancer Society website. Updated June 26, 2024. Accessed November 3, 2025. <https://www.cancer.org/cancer/managing-cancer/side-effects/eating-problems/nausea-and-vomiting/managing.html>. 9. National Cancer Institute. Cancer fatigue. National Cancer Institute website. Updated September 20, 2024. Accessed August 27, 2025. <https://www.cancer.gov/about-cancer/treatment/side-effects/fatigue>. 10. Cleveland Clinic. Cancer fatigue. Cleveland Clinic website. Updated April 16, 2024. Accessed November 3, 2025. <https://my.clevelandclinic.org/health/diseases/5230-cancer-fatigue>.

 Bristol Myers Squibb®

© 2025 Mirati Therapeutics, Inc., a Bristol Myers Squibb company.
KRAZATI and the related logo are registered trademarks of Mirati Therapeutics, Inc.
1914-US-2500116 11/25

Round

M20

PRINT

Subsidiary:			
Client:	BMS		
Product:	Krazati		
Job #:	42659-25XKRAZAT		
Name:	DTC Patient FAQs Update (Print)		
Path:	/Users/hencizo/Library/CloudStorage/Box-Box/DEPARTMENT_CREATIVE_STUDIO/BMS/Krazati_KRAZI/42659-25XKRAZAT_FAQsUpdate/Print/42659-25XKRAZATFAQs_Update_Print_M20		
File Name:	42659-25XKRAZAT FAQs_Update_Print_M20	Date:	November24,20252:46PM

Bleed:	.25"
Trim:	8" (w) x 8' (h)
Safety:	.25"
Inks:	6/6 (Process + PMS 2270 + PMS 2728)

Studio Artist	Hank E
Cross Check	
Art Lead	Mirian • PRE-PRO: Gerard G
Studio Lead	Anthony • PRE-PRO: Donna K
Production Manager	Mark • PRE-PRO: Mark R
Project Manager	Kaitlyn • PRE-PRO: Kaitlin G

Notes:

Pick up file: 42659-DTC_FAQ_Update_Print_V17

Use Connect Fonts